

Jet quenching and Direct Photon Production

Fu-Ming Liu,¹ Tetsufumi Hirano,² Klaus Werner,³ and Yan Zhu¹

¹*Institute of Particle Physics, Central China Normal University, Wuhan, China*

²*Department of Physics, The University of Tokyo, 113-0033, Japan*

³*Laboratoire SUBATECH, University of Nantes - IN2P3/CNRS - Ecole des Mines, Nantes, France*

(Dated: November 11, 2008)

Jet quenching effect has been investigated in the direct photon production, based on a realistic data-constrained (3+1) dimensional hydrodynamic description of the expanding hot and dense matter, a reasonable treatment of the propagation of partons and their energy loss in the fluid, and a systematic study of the main sources of direct photons. Our resultant p_t spectra agree with recent PHENIX data in a broad p_t range. Parton energy loss in the plasma eventually effect significantly direct photon production from fragmentation and jet photon conversion, similar to hadron suppression in central heavy ion collisions. But this only causes about 40% decrease in the total production of direct photons, due to the mixture with other direct photon sources.

PACS numbers:

I. INTRODUCTION

The formation and observation of a quark-gluon plasma (QGP) in heavy ion collisions are important goals of modern nuclear physics [1, 2]. Suppression of high p_t hadron yields [3] is one of the most important features observed at the Relativistic Heavy Ion Collider (RHIC). Theoretically this is attributed to the interaction between jets (hard partons) and the bulk matter [4, 5, 6, 7]. This offers us an excellent opportunity to study the interaction of partons inside the system and, consequently, properties of the matter under extreme conditions.

However the observed centrality-dependence of hadronic jet quenching doesn't show up in direct photon production[8]. In this paper, we investigate the role of jet quenching on direct photon production. For this purpose, we need a careful consideration of three components: 1) the space-time evolution of the hot dense matter or the bulk matter; 2) hard parton production and propagation in the bulk and 3) all sources of direct photons.

The space-time evolution of the hot dense matter is achieved by using three dimensional (3D) hydrodynamic simulations of bulk matter [10, 11] which have already been tested against a vast body of low p_t hadron data at RHIC.

The interaction between hard parton and the bulk so that energy loss of the hard parton is formulated via the BDMPs framework [7] and tested on pion suppression at various centralities.

Four significant sources are considered in this paper: leading order contribution in Primordial NN scattering, thermal contribution, contribution from jet-photon conversion and from jet fragmentation. There are possible sources ignored at present study: The medium-

induced photon radiation is expected to contribute at low transverse momentum region where thermal contribution dominants. The interaction involving non-equilibrated soft partons may also produce direct photons, but the preequilibrium time interval is much shorter than the life time of the equilibrated matter (~ 20 fm/c).

Therefore energy loss of hard parton in the bulk will directly effect two direct photon sources, jet-photon conversion and from jet fragmentation. We will see the role of jet quenching on direct photon production clearly via investigating those production in Au-Au collisions at different centralities where various sizes of hot dense matter are formed. Certainly different sizes of bulks will also contribute thermal photon production differently which will not be discussed in detail in this paper.

The paper is organized as follows: In Sec. II, we first give a brief review on the space-time evolution of the hot matter created in Au+Au collisions at different centralities based on a (3+1)-dimensional ideal hydrodynamical calculation. In Sec. III, we discuss parton energy loss in the QGP. We investigate neutral pion production in the high p_t region in order to fix the parameters of the energy loss scheme. We discuss sequently the contributions from various sources to direct photon p_t spectra in Sec. IV. We show our results and compare them with recent experimental data in Sec. V. Section VI is devoted to conclusion of the present study.

II. SPACE-TIME EVOLUTION OF THE HOT AND DENSE MATTER

In our calculation, fully three-dimensional (3D) ideal hydrodynamics [10, 11] is employed to describe the space-time evolution of the hot and dense matter created in

Au+Au collisions at RHIC energy at various centralities. We solve the equations of energy-momentum conservation

$$\partial_\mu T^{\mu\nu} = 0 \quad (1)$$

in full 3D space (τ, x, y, η) under the assumption that the local thermal equilibrium is reached (maintained) at (after) an initial time $\tau_0 = 0.6 \text{ fm}/c$. Here τ and η are the proper time and the space-time rapidity, respectively. x and y are transverse coordinates. In the transverse plane, the centers of two colliding nuclei are located at $(x, y) = (b/2, 0)$ and $(-b/2, 0)$ before the collision at an impact parameter b . Ideal hydrodynamics is characterized by the energy-momentum tensor,

$$T^{\mu\nu} = (e + P)u^\mu u^\nu - Pg^{\mu\nu}, \quad (2)$$

where e , P , and u^μ are energy density, pressure, and local four velocity, respectively. We neglect the finite net-baryon density which is small near the mid-rapidity at RHIC. For the high temperature ($T > T_c = 170 \text{ MeV}$) QGP phase we use the equation of state (EOS) of massless non-interacting parton gas (u , d , s quarks and gluons) with a bag pressure B :

$$p = \frac{1}{3}(e - 4B). \quad (3)$$

The bag constant is tuned to be $B^{\frac{1}{4}} = 247.19 \text{ MeV}$ to match pressure of the QGP phase to that of a hadron resonance gas at critical temperature $T_c = 170 \text{ MeV}$. A hadron resonance gas model at $T < T_c$ includes all hadrons up to the mass of the $\Delta(1232)$ resonance. Our hadron resonance gas EOS implements chemical freeze-out at $T_{\text{ch}} = T_c = 170 \text{ MeV}$, as observed in collisions at RHIC [12].

We assume that, at $\tau_0 = 0.6 \text{ fm}/c$, the initial entropy distributions is proportional to a linear combination of the number density of participants (85%) and binary collisions (15%) [13]. Centrality dependence of charged particle multiplicity observed by PHOBOS [14] has been well reproduced by full 3D hydrodynamics simulations with the above setups [13]. In the following calculations, hydrodynamic outputs at representative impact parameters $b = 3.2, 5.5, 7.2, 8.5, 9.7$, and 10.8 fm are chosen for 0-10%, 10-20%, $\dots$, 50-60% centrality, respectively.

For convenience of the following calculations, we introduce $f_{\text{QGP}}(\tau, x, y, \eta)$ as the fraction of the QGP phase in a fluid element. It is obvious that $f_{\text{QGP}} = 1$ (0) in the QGP (hadronic) phase. In the mixed phase, the fraction of the QGP is calculated via

$$f_{\text{QGP}} e_{\text{QGP}} + (1 - f_{\text{QGP}}) e_{\text{had}} = e(\tau, x, y, \eta)$$

with e_{QGP} and e_{had} being the energy densities of the QGP phase and the hadron phase at $T = T_c$, respectively.

III. PARTON ENERGY LOSS IN A PLASMA

Hard parton production from primordial nucleus-nucleus scattering is calculated as [15]

$$\frac{dN^{AB \rightarrow \text{jet}}}{dy d^2 p_t} = K T_{AB}(b) \sum_{abcd} \int dx_a dx_b G_{a/A}(x_a, M^2) \times G_{b/B}(x_b, M^2) \frac{\hat{s}}{\pi} \frac{d\sigma}{d\hat{t}}(ab \rightarrow cd) \delta(\hat{s} + \hat{t} + \hat{u}) \quad (4)$$

where $T_{AB}(b)$ is the nuclear overlapping function at an impact parameter b for each centrality, $G_{a/A}(x_a, M^2)$ and $G_{b/B}(x_b, M^2)$ are parton distribution functions in nuclei A and B . We take MRST 2001 LO parton distributions in proton [16]. $K = 2$ is chosen to take into account higher order contributions. These parameters are chosen as to reproduce high p_t pion data in pp collisions at RHIC, which will be discussed later. Nuclear shadowing effect and EMC effect are taken into account through EKS98 scale dependent nuclear ratios $R_a^{\text{EKS}}(x, A)$ [17]. Isospin of a nucleus with mass A , neutron number N , and proton number Z is corrected as follows:

$$G_{a/A}(x) = \left[\frac{N}{A} G_{a/N}(x) + \frac{Z}{A} G_{a/P}(x) \right] R_a^{\text{EKS}}(x, A). \quad (5)$$

The isospin mixture and nuclear shadowing eventually cause a decrease of nuclear modification at high p_t region.

We assume that all jets are produced at $\tau = 1/Q \approx 0$ with the phase space distribution

$$f_0(\vec{p}, \vec{r}) \propto \frac{dN}{d^3 p} T_A \left(x - \frac{b}{2}, y \right) T_B \left(x + \frac{b}{2}, y \right) \delta(z) \quad (6)$$

where $\vec{r} = (x, y, z)$ is the coordinate of a jet, b is the impact parameter, and T_A and T_B are thickness functions of nuclei A and B .

When hard partons propagate in a hot dense matter, they may lose energy in the medium via radiation or elastic collisions with thermal partons [4, 5, 6, 7]. This is a quite hot issue and several different schemes have been employed for jet energy loss in QGP phase while the study of hard parton energy loss in hadronic phase is not so popular. In this paper the BDMPS schemes is employed which considers the energy loss of hard parton in QGP phase via radiation. A free parameter D is introduced by us to take into account of energy loss via other modes.

The total energy loss along the trajectory of a hard parton is calculated as

$$\Delta E(i, \vec{p}_0, \vec{r}_0) = D \int_{\tau_0}^{\infty} d\tau \epsilon(i, \tau, \mathbf{x}(\tau)) \theta(f_{\text{QGP}}(\tau, \mathbf{x}(\tau))), \quad (7)$$

where $\mathbf{x}(\tau)$ is the trajectory of parton, $f_{\text{QGP}}(\tau, \mathbf{x}(\tau))$ is the fraction of the QGP phase at a position $(\tau, \mathbf{x}(\tau))$, and θ is a step function, which gives $\theta(f_{\text{QGP}})$ equal unity in the QGP and the mixed phases and zero in the hadron

phase. In the mixed phase, the thermal parton density will be diluted with the factor f_{QGP} , the fraction of the QGP phase. Here D is an adjustable parameter, $\epsilon(i, \tau, \mathbf{x}(\tau))$ is the energy loss per unit distance for a parton i at a position $(\tau, \mathbf{x}(\tau))$, given as [6]

$$\epsilon(i, \tau, \mathbf{x}(\tau)) = \alpha_s \sqrt{\mu^2 E^* / \lambda_i}.$$

Note that the above quantities, *i.e.*, temperature T , fluid velocity u_μ , parton densities ρ_i and, in turn, mean free path λ_i , depend on the location of the parton $\mathbf{x}(\tau)$ and can be obtained from full 3D hydrodynamics simulations discussed in the previous section.

We first discuss pion production in proton-proton collisions. We calculate neutral π -meson production assuming pQCD factorization, Eq. (4),

$$\frac{dN_{pp}^{\pi^0}}{dyd^2p_t} = \sum_{c=g,q_i} \int dz_c \frac{dN^{pp \rightarrow c}}{dyd^2p_t^c} \frac{1}{z_c^2} D_{\pi^0/c}^0(z_c, Q^2), \quad (8)$$

where $D_{\pi^0/c}^0(z_c, Q^2)$ is π^0 fragmentation functions parameterized by Kniehl *et al.* [18]. In the high p_t region where the pQCD is expected to work, we reasonably reproduce the experimental data with the above setup with $K = 2$ and $M = Q = p_t$. We use the p_t spectrum as a reference spectrum in the following calculations.

The effect of parton energy loss is taken into account through the medium modified fragmentation function [19] $D_{\pi^0/c}(z_c, Q^2, \Delta E_c)$. With a common value of the parameter $D = 1.5$, we can reasonably reproduce the PHENIX data [9] on π^0 jet quenching in the high p_t region at all centralities simultaneously. In the following photon calculations, we always use the BDMPS energy loss formula (7) with $D = 1.5$.

IV. THE DIFFERENT SOURCES OF DIRECT PHOTON PRODUCTION

Possible sources of direct photons are considered in the following.

Leading order contribution in Primordial NN scattering. The direct photon production via Compton scattering and quark-antiquark annihilation can be calculated in perturbation theory using the conventional parton distribution functions and the factorization hypothesis. Higher order contributions are considered as a part of the so-called jet fragmentation contribution and jet photon conversion which will be discussed separately. Similar to Eq. (4), the leading order contribution to direct photon production in nucleus-nucleus collisions reads

$$\begin{aligned} \frac{dN^{AB \rightarrow \gamma}}{dyd^2p_t} &= T_{AB}(b) \sum_{ab} \int dx_a dx_b G_{a/A}(x_a, M^2) \\ &\times G_{b/B}(x_b, M^2) \frac{\hat{s}}{\pi} \frac{d\sigma}{d\hat{t}}(ab \rightarrow \gamma + X) \delta(\hat{s} + \hat{t} + \hat{u}) \end{aligned} \quad (9)$$

where the elementary processes $ab \rightarrow \gamma + X$ are Compton scattering $qg \rightarrow \gamma q$ and annihilation $q\bar{q} \rightarrow g\gamma$.

Thermal production. In high energy nuclear collisions, the density of secondary partons is so high that the quarks and gluons rescatter and eventually thermalize to form a bubble of hot QGP. Thermal photons can be produced via interaction between bulk particles during the whole history of the evolution of hot matter. The interaction of bulk particles and the bulk properties can be reflected by the emission rate of direct photons in the QGP phase[21] and in the hadronic phase[22], noted as $\Gamma = Ed^3R/d^3p$, where R is the number of photons emitted from a medium per unit space-time volume with temperature T . Total yields of thermal photons can be obtained by summing the emission rate over the space-time volume as

$$\frac{dN^{\text{thermal}}}{dyd^2p_t} = \int d^4x \Gamma(p^\mu u_\mu, T) \quad (10)$$

with $d^4x = \tau d\tau dx dy d\eta$, p^μ the photon's four momentum in the laboratory frame and u_μ a local fluid velocity.

Jet fragmentation. At any stages of the evolution of a jet (final state parton emission), there is a possibility of emitting photons. The existence of a QGP affects the results because energetic partons lose their energy prior to fragmentation. In this work, we assume fragmentation of partons only outside the plasma, which is similar to high p_t hadron production from jet fragmentation.

In pp collisions, there is no formation of QGP and no energy loss of hard parton in QGP. Higher order contributions to direct photon production can be considered as jet fragmentation and be calculated as

$$\frac{dN_{pp}^{\text{frag}}}{dyd^2p_t} = \sum_{c=g,q_i} \int dz_c \frac{dN^{pp \rightarrow c}}{dyd^2p_t^c} \frac{1}{z_c^2} D_{\gamma/c}^0(z_c, Q^2), \quad (11)$$

with photon fragmentation functions $D_{\gamma/c}^0(z, Q^2)$ being the probability for obtaining a photon from a parton c which carries a fraction z of the parton's momentum. So $p_t^c = p_t/z_c$ is the transverse momentum carried by the parton c before fragmentation and $d^3p/E = z_c^2 d^3p^c/E^c$. The effective fragmentation functions for obtaining photons from partons can be calculated perturbatively. We use the parameterized solutions by Owens [15].

In case of heavy ion collisions where a QGP is formed, energy loss of hard partons in a plasma should be taken into account. This can be done via modified fragmentation functions [19]

$$\begin{aligned} D_{\gamma/c}(z_c, Q^2, \Delta E_c) &= \left(1 - e^{-\frac{L}{\lambda_c}}\right) \left[\frac{z'_c}{z_c} D_{\gamma/c}^0(z'_c, Q^2) + \frac{L}{\lambda_c} \frac{z'_g}{z_c} D_{\gamma/g}^0(z'_g, Q^2) \right] \\ &+ e^{-\frac{L}{\lambda_c}} D_{\gamma/c}^0(z_c, Q^2), \end{aligned} \quad (12)$$

with $z'_c = p_t/(p_t^c - \Delta E_c)$ and $z'_g = (L/\lambda_c) p_t/\Delta E_c$ being the rescaled momentum fractions carried by the parton c and the emitted gluons before fragmentation. λ_c is mean free path of the parton c in the plasma, L is the path length of each parton traversing the plasma. Thus,

in heavy ion collisions, contributions from fragmentation become

$$\frac{dN_{AB}^{\text{frag}}}{dyd^2p_t} = \sum_{c=g,q_i} \int dz_c \frac{dN^{AB \rightarrow c}}{dyd^2p_t^c} \frac{1}{z_c^2} D_{\gamma/c}(z_c, Q^2, \Delta E_c). \quad (13)$$

Jet photon conversion with jet energy loss. When hard partons propagate in a plasma, they also collide with thermal partons and produce direct photons via Compton process and the quark-antiquark annihilation. We call this process jet-photon conversion, since it is a conversion of a jet into a photon with almost the same momentum as the one of originating jet parton. Contribution from the jet-photon conversion is calculated by integration of conversion rate over the space-time evolution of the hot and dense matter in the QGP phase

$$\frac{dN^{\text{jpc}}}{dyd^2p_t} = \int \Gamma^{\text{jpc}}(E^*, T) f_{\text{QGP}}(x, y, \eta, \tau) d^4x. \quad (14)$$

The photon production rate by annihilation and Compton scattering of hard partons in the medium [23, 24]

$$\Gamma^{\text{jpc}}(E^*, T) = \frac{\alpha \alpha_s}{4\pi^2} \sum_q e_q^2 f_q(\vec{p}, x) T^2 \left[\ln \frac{4E_\gamma^* T}{m_{\text{th}}^2} - C \right] \quad (15)$$

where E^* is the photon energy in the local rest frame, $C = 2.323$, $m_{\text{th}}^2 = g^2 T^2/6$, and the strong coupling $\alpha_s = g^2/4\pi$ being temperature dependent as in Eq. (??). α is the electromagnetic couplings, e_q and $f_q(\vec{p}, x)$ are the electric charge and the phase-space density of a hard parton of flavor q . The phase space distribution of hard partons at τ is obtained by considering parton energy loss as

$$\begin{aligned} f(\vec{p}, x) &= f(\vec{p}, \vec{r}, \tau) \\ &= \int d^3p_0 f_0(\vec{p}_0, \vec{r} - \vec{v}t) \delta(\vec{p}_0 - \vec{p} - \vec{v}\Delta E) \end{aligned}$$

where $f_0(\vec{p}, \vec{r})$ is the phase space distribution at $\tau = 0$ described in Eqs. (6). The δ -function expression reflects the energy loss of a parton moving along a straight line trajectory with $\vec{v} \equiv \vec{p}/E = \vec{p}_0/E_0$. ΔE is the energy loss from τ_0 to τ and calculated similar to Eq. (7) but replacing the upper limit of integral ∞ with τ .

V. RESULTS AND DISCUSSION

In Fig. 1, the calculated p_t spectra of direct photons in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV at centrality 0-20% and 20-40% are compared to PHENIX data [25, 26]. The PHENIX data are reproduced within our multi-component model remarkably well.

The nuclear modification factor R_{AA} is obtained by dividing a p_t spectrum in nucleus-nucleus collisions by the N_{coll} -scaled p_t spectrum in pp collisions. The calculation of direct photons in pp collisions includes the leading order contribution plus fragmentation contribution.

In high p_t regions, our result agrees with the PHENIX data [25], [26] reasonably well: So we use it to calculate nuclear modification. It also provides a baseline calculation with the LO contribution and fragmentation contribution in Au+Au collisions.

Figure 2 shows how the nuclear modification factor for direct photons, R_{AA} , depends on centrality and on energy loss. Data for 0-10% centrality are taken from Refs. [27] and [28]. Figure 2(a) shows centrality dependence of R_{AA} compared to the PHENIX data. The three curves are respectively 0-10% (dotted line), 20-30% (solid line), and 40-50% (dash-dotted line).

R_{AA} has a weak centrality dependence at high p_t region. This result is consistent with the observed phenomenon [27] that the p_t -integrated (for $p_t > 6$ GeV/c) R_{AA} of direct photons is almost independent of collision centrality. Does this imply a very weak effect from jet quenching? Figure 2(b) answers this question (here for the most central collisions): Comparing calculations with (dotted line) and without energy loss (solid line), one finds a difference of up to 40%. So the effect of parton energy loss is quite visible in the p_t range between 4 GeV/c and more than 20 GeV/c. If we would do the R_{AA} calculations without energy loss, the difference between central and semiperipheral collisions would be about 20%, whereas the complete calculation gives the same result for all centralities, within 5%. We have to admit that we talk about small effects, requiring experimental data with relative errors of less than 5% to observe the effects.

Parton energy loss in the plasma suppresses the fragmentation contributions and jet-photon conversion. So we study the ratios of the contribution with energy loss to the one without energy loss, as shown in Fig. 3 ((a) for fragmentation and (b) for jet-photon conversion). Energy loss in the plasma depends on the path length of the hard parton inside the plasma, which turns out to depend on the collision centrality. We do see a similar centrality dependence of the suppression for π^0 (jet quenching effect) in fragmentation contributions and jet-photon conversion.

VI. CONCLUSION

We calculated the centrality-dependence of p_t spectra for direct photons in Au+Au collisions at the RHIC energy, based on a realistic data-constrained (3+1)-D hydrodynamic description of the expanding hot and dense matter, a reasonable treatment of propagation of partons and their energy loss, and a systematic consideration of main sources of direct photons. In this study, four main sources are considered, namely, leading order (LO) contribution from primordial elementary scatterings, thermal radiation from the fluids, fragmentation from hard partons, and jet photon conversion (JPC). Similar work [29] has been done before the appearance of the most recent data [26]. Our results agree nicely with the recent low p_t data.




Figure 1: (Color Online) Direct photon production in Au+Au collisions at centrality 0-20% and 20-40%. PHENIX data are shown as open circles [25] and filled circles [26].




Figure 2: (Color Online) The nuclear modification factor of direct photons in Au+Au collisions R_{AA} . Data for 0-10% centrality are from Ref. [27] and Ref. [28]. (a): R_{AA} at centrality 0-10% (dotted line), 20-30% (solid line), and 40-50% (dash-dotted line) respectively. (b): R_{AA} at 0-10% centrality with energy loss (dotted line) and without energy loss (solid line).

The role of jet quenching in the high p_t region of direct photons production has been checked via fragmentation photons and jet photon conversion sources. For these two sources, the suppression of the photon rate due to parton energy loss is significant in central Au+Au collisions, and becomes less important towards peripheral collisions, similar to the suppression for meson production. Since experimentally one may separate isolated photons (LO+JPC) and associate photons (fragmentation photons), our prediction may be tested in the future.

Considering the total yields of direct photons, the contribution from fragmentation and conversion are small, contributing between 5% and 10%. However, parton energy loss plays nevertheless an important role: Without it, these second order effects would contribute significantly. Without jet quenching, the nuclear modification

factors R_{AA} would depend visibly on the centrality of the collisions. A strong energy loss is actually necessary to get the centrality scaling of R_{AA} in our calculation – a centrality scaling which has been observed by the PHENIX collaboration. In this sense, properties of the bulk matter affect the photon yields at intermediate values of p_t , via the parton energy loss.

Acknowledgments

This work is supported by the Natural Science Foundation of China under the project No. 10505010 and MOE of China under project No. IRT0624. The work of T.H. was partly supported by Grant-in-Aid for Scientific Research No. 19740130.

[1] Proceedings of Quark Matter 2006, J. Phys. G: Nucl. Part. Phys. **34** (2007).
 [2] J. W. Harris and B. Muller, Ann. Rev. Nucl. Part. Sci. **46**, 71 (1996).

[3] K. Adcox *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **88**, 022301 (2002); S.S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **91**, 072301 (2003); C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **89**, 202301




Figure 3: (Color Online) The ratio of the contribution with energy loss to the one without, in fragmentation (a) and jet-photon conversion (b).

- (2002); C. Adler *et al.* [STAR Collaboration], Phys. Rev. Lett. **90**, 082302 (2003).
- [4] J. D. Bjorken, FERMILAB-PUB-82-059-THY.
- [5] M. Gyulassy and M. Plumer, Phys. Lett. B **243**, 432 (1990); X.N. Wang and M. Gyulassy, Phys. Rev. Lett. **68**, 1480 (1992).
- [6] R. Baier, Y.L. Dokshitzer, A.H. Mueller, S. Peigne, and D. Schiff, Nucl. Phys. B **483**, 291 (1997); Nucl. Phys. B **484**, 265 (1997).
- [7] R. Baier, D. Schiff, B.G. Zakharov, Ann. Rev. Nucl. Part. Sci. **50**, 37 (2000); U. Wiedemann, Nucl. Phys. B **588**, 303 (2000).
- [8] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **96**, 202301 (2006) [arXiv:nucl-ex/0601037].
- [9] A. Adare *et al.* [PHENIX Collaboration], arXiv:0801.4020 [nucl-ex].
- [10] T. Hirano, Phys. Rev. C **65**, 011901 (2001).
- [11] T. Hirano and K. Tsuda, Phys. Rev. C **66**, 054905 (2002).
- [12] P. Braun-Munzinger, D. Magestro, K. Redlich, and J. Stachel, Phys. Lett. B **518**, 41 (2001).
- [13] T. Hirano, U. Heinz, D. Kharzeev, R. Lacey, and Y. Nara, Phys. Lett. B **636**, 299 (2006); J. Phys. G **34**, S879 (2007); Phys. Rev. C **77**, 044909 (2008).
- [14] B.B. Back *et al.* [PHOBOS Collaboration], Phys. Rev. C **65**, 061901 (2002).
- [15] J.F. Owens, Rev. Mod. Phys. **59**, 465 (1987).
- [16] A. D. Martin, R. G. Roberts, W. J. Stirling, and R. S. Thorne, Eur. Phys. J. C **23**, 73 (2002).
- [17] K.J. Eskola, V.J. Kolhinen and C.A. Salgado, Eur. Phys. J. C **9**, 61 (1999); K.J. Eskola, V.J. Kolhinen and P.V. Ruuskanen, Nucl. Phys. B **535**, 351 (1998).
- [18] B.A. Kniehl, G. Kramer, and B. Potter, Nucl. Phys. B **597** 337 (2001).
- [19] X.N. Wang, Phys. Lett. B **595**, 165 (2004); **570**, 299 (2004).
- [20] R. Baier, H. Nakkagawa, A. Niegawa, and K. Redlich, Z. Phys. C **53**, 433 (1992).
- [21] P. Arnold, G. D. Moore, and L. G. Yaffe, J. High Energy Phys. **0111**, 057 (2001); J. High Energy Phys. **0112**, 9 (2001).
- [22] S. Turbide, R. Rapp and C. Gale, Phys. Rev. C **69**, 014903 (2004); C. Song, Phys. Rev. C **47**, 2861 (1993).
- [23] C.Y. Wang, Introduction to high-energy heavy ion collisions (World Scientific Singapore, 1994).
- [24] R.J. Fries, B. Muller, and D.K. Srivastava, Phys. Rev. Lett. **90**, 132301 (2003); R. J. Fries, B. Muller and D. K. Srivastava, Phys. Rev. C **72**, 041902 (2005).
- [25] S. S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **98**, 012002 (2007).
- [26] A. Adare *et al.* [PHENIX Collaboration], arXiv:0804.4168 [nucl-ex].
- [27] S.S. Adler *et al.* [PHENIX Collaboration], Phys. Rev. Lett. **94**, 232301 (2005).
- [28] Tadaaki Isobe *et al.* [PHENIX Collaboration] J. Phys. G **34**, S1015 (2007).
- [29] S. Turbide, C. Gale, E. Frodermann and U. Heinz, Phys. Rev. C **77**, 024909 (2008).